

HEALTH INDICATOR

2007/08

ARHS

HARAMAYA UNIVERSITY



Age ranged between 15 years and 19 years and the median age was 17 years

No	Indicators	Percent
1	Adolescents not able to read and write in the study population	38.7
2	Male adolescents not able to read and write	28.3
3	Female adolescents not able to read and write	49.8
4	Adolescents who were married	9.4
5	The prevalence of early marriage in the study population	9.4
6	The prevalence of early marriage for males	4.9
7	The prevalence of early marriage in the females population	14.2
8	Adolescents ever had received information on ARH	3
9	Adolescents who had known the availability of ARH service in their locality reported visiting the service providing unit	12.5
10	Adolescents who were sexually active	12.9
11	Males adolescents were sexually active	5.7
12	Females adolescents were sexually active	7.2
13	Adolescents were knowledgeable about condom	32.9
14	Male adolescents were knowledgeable about condom	45.6
15	Female adolescents were knowledgeable about condom	19.4
16	Sexually active adolescents were knowledgeable about condom	42.1
17	Sexually active adolescents ever using condoms	14
18	Sexually active male adolescents ever using condoms	21.3
19	Sexually active female adolescents ever using condoms	8.3
20	Married female adolescents using modern contraceptive methods	3.5
21	Unmarried female adolescents using modern contraceptive methods	60
22	Adolescents utilized VCT	1.6
23	Male adolescents utilized VCT	2.3
24	Female adolescents utilized VCT	0.7
25	Adolescents discussed on RH issues with their parents	13.0
26	Adolescents male discussed on RH issues with their parents	60.4%
27	Adolescents female discussed on RH issues with their parents	39.6%



College of Health & Medical Sciences, KDS-HRC

✉ 235, Harar, Ethiopia

Email: kds_hrc@yahoo.com

☎ 00251-256-66-6143

HEALTH INDICATOR

2007/08

HARAMAYA UNIVERSITY



No	Indicators	Percent
28	Know ledge of adolescent reproductive health (ARH) among adolescents	20.3
29	Knowledge of menarche among female adolescents	17.0
30	Knowledge of timing of semen ejaculation among male adolescents	36.1
31	Knowledge of method of preventing pregnancy	38.4
32	Prevalence of sexual coercion among females adolescents	7.0
33	Magnitude of early sexual debut	13.7
34	Prevalence of multiple life time sexual partners.	17.8
35	Ever having communication about sexuality with parents among adolescents	25.6
36	Ever had known about condom	31.2
37	Ever use of condom	6.1
38	Know the disease that can be transmitted sexually	37.5
39	Perceived risk of acquiring sexually transmitted infections	11.0
40	Have ever heard of HIV/AIDS	51.6
41	Nont risk perception of acquiring HIV/ AIDS	71.6

No	Indicators	Value
1	The mean age at the first sexual debut	16.62 Years



College of Health & Medical Sciences, KDS-HRC

✉ 235, Harar, Ethiopia

Email: kds_hrc@yahoo.com

☎ 00251-256-66-6143